

CASE REPORT

Open Access



Orthodontic Extrusion with crown restoration in the management of anterior teeth crown-root fracture: A case report

Wang Jing¹, Huo Na¹, Cai Chuan¹, Xu Lulu^{1*}

Abstract

Crown-root fracture is a rare type of dental traumatic injury. Management of traumatized teeth sometimes requires multidisciplinary approach to obtain better esthetic results as well as for conservation of tooth structure. This article describes the management of a case of oblique crown root fracture of anterior teeth. After removal of the coronal fragment, the roots were filled with calcium hydroxide. Orthodontic extrusion was initiated for 4months and 3mm extrusion was achieved. Ceramic crown was cemented on anterior fracture teeth. The periodontal condition appeared to be healthy and good esthetic was maintained during the follow up appointment.

Key words: Fracture, Orthodontic Extrusion, Anterior teeth, Ceramic crown

*Correspondence:

Xu Lulu
Department of Stomatology,
The First Medical Center of PLA General Hospital,
28 Fuxing Road, Haidian District, Beijing
China

¹Department of Stomatology, The First
Medical Center of PLA General
Hospital, 28 Fuxing Road, Haidian
District, Beijing, China

Received Date: June 11, 2026; **Accepted date:** June 22, 2026; **Published Date:** July 07, 2026

Citation: Xu Lulu; Arch Med Clin Case Stud; "Orthodontic Extrusion with crown restoration in the management of anterior teeth crown-root fracture: A case report". 2026; 1(1): 132



Copyright: © Xu Lulu 2026. This is an open-access article distributed under the terms of the Creative Commons Attribution License.

INTRODUCTION

Crown-root fracture is a rare type of dental traumatic injury, affecting 0.5% to 7% of permanent teeth¹. In this type of injury, the wound involves damage to the pulp, cementum, dentin, and periodontal ligament. Depending on the severity of the injury, multiple or single fractures of the root can be observed. Management of traumatized teeth sometimes requires multidisciplinary approach to obtain better esthetic results as well as for conservation of tooth structure². Exposure of the fractured margin into the oral cavity is desirable. This may be achieved by³: (i) crown lengthening, (ii) surgical extrusion, (iii) orthodontic extrusion. This case report presents management of a cervical root fracture using orthodontic extrusion and a final restoration with ceramic crown.

Case Report:

A 22-year-old man reported to the Department of Orthodontic Dentistry with a complaint of damaged anterior teeth. History revealed that the patient had an injury to the front tooth

gingivally on the mesiobuccal and mesiolingual aspect. Periapical radiograph revealed that the fracture line was extending up to the level of alveolar crest. The root canal treatment of fractured teeth was finished (Fig1B).

Since the fracture line extended up to the alveolar bone crest, it was possible to initiate extrusion of the root so as to facilitate placement of a coronal restoration and long-term attachment health. So orthodontic extrusion was possible. Lingual button was bonded on fracture side of the anterior teeth (Fig2A). The patient was revisioned once a month and orthodontic traction was taken. Removable appliances were used for lower arch to tract anterior teeth with help of the buttons. A small amount of both labial and palatal mucous recontouring was done using electrotome method to allow for proper finishing of the crown margins (Fig2B). Interactive traction was used for 4 months, and 3 mm extrusion was achieved (Fig2C, 2D). The temporary crown was used for 6 months (Fig3A). Finally ceramic crown was cemented




Fig1A: Intraoral examination revealed anterior teeth crown-root fractured, the coronal fragment was extracted

Fig1B: Periapical radiograph revealed that the fracture line was extending up to the level of alveolar crest. The root canal treatment of fractured teeth was finished

region 1 month ago, due to a fall. Intraoral examination revealed anterior teeth crown-root fractured, the coronal fragment was extracted (Fig1A). The fracture line was located sub

on 12 11 and 21 (Fig3B). During the three month follow up appointment, the periodontium appeared to be healthy and good esthetic was maintained.




Fig2A: Lingual button was bonded on fracture side of the anterior teeth.

Fig2B: A small amount of both labial and palatal mucosal recontouring was done using electrosurgical method to allow for proper finishing of the crown margins

Fig2C& 2D: Interactive traction was used for 4 months and 3 mm extrusion was achieved




Fig3A: The temporary crown was used for 6 months

Fig3B: Ceramic crown was cemented on 12 11 and 21

DICUSSION

Orthodontic forced eruption was first described by Heithersay as coronal orthodontic movement achieved by applying continuous orthodontic force for the purpose of preserving soft tissues and gaining vertical bone height. Orthodontic extrusion is a safe, minimally invasive and highly predictable treatment. It shows several advantages if compared to

surgical crown lengthening⁴. First, tooth structure and periodontal support maintenance is crucial. As previously mentioned, it is required to extend bone resection on the adjacent teeth to harmonize the gingival morphology in surgical crown lengthening. This causes bone loss and a possible damage of the periodontal support as well as a worsening of the crown/root ratio of these teeth. Conversely, orthodontic extrusion can just

cause tooth displacement or even increase the volume of dental support tissues, which is particularly advantageous for implant purposes⁵.

Orthodontic extrusion can be obtained using different orthodontic strategies⁶: fixed appliances, removable appliances, temporary anchorage devices such as mini-screw. Different treatment choice variables must be considered, such as patient preference, oral hygiene, availability of an appropriate orthodontic anchorage and amount of dental crown available. As a general rule, the less periodontal support, the less force needed for extrusion. An accelerated movement rate may result in a risk of tissue damage and ankylosis. A sufficient crown-root-ratio (ideally less than 1:1), healthy gingiva, and osseous structures without any pathological signs are required for successful extrusion and favorable esthetic outcomes in crown-root fractured teeth. The current knowledge suggests the application of light and constant forces of 15 g for the anterior teeth and 50 g for posterior teeth are optimal to extrude a tooth for implant site enhancement⁷. After successful application of orthodontic extrusion, the stabilization phase takes place. Resin crown was used as temporary crown for 6 months. This phase is important to allow for proper reorganization of the soft tissue and remodeling of the bone in the so-called “osteoid formation” phase, thus discouraging relapse. It is preferable to choose orthodontic extrusion if a highly predictable treatment is requested, if an orthodontic device is already present and if it is necessary to preserve tooth vitality or treat teeth non-compatible with an atraumatic extraction. Orthodontic movements, especially if performed on traumatized teeth or through the application of heavy forces, can cause root resorption⁸. However, it can be considered a rare event when performing extrusive movement. Moreover, there was no correlation between the degree of extrusion and the occurrence of root resorption.

CONCLUSION

Using a combination of orthodontic extrusion and crown restoration may be considered for treatment of crown root fractures because:

- 1) Esthetics is maintained until the final restoration placement by highly cost effective.
- 2) Excellent esthetics and periodontal health is maintained if the final restoration is made after orthodontic extrusion.
- 3) It is a simple and safe method and hence more acceptable to the patient.

The prospects of this treatment modality for crown root fractures of posterior teeth needs to be investigated.

REFERENCES

1. Martina Cordaro, Edoardo Staderini, Ferruccio Torsello, Nicola Maria Grande, Matteo Turchi, Massimo Cordaro. Orthodontic Extrusion vs. Surgical Extrusion to Rehabilitate Severely Damaged Teeth: A Literature Review. *Int J Environ Res Public Health*. 2021;18, 9530.
2. Abdulaziz Alsahhaf BDSa, Wael Att, Med Dent. Orthodontic extrusion for pre-implant site enhancement: Principles and clinical guidelines. *Journal of prosthodontic research*. 60(2016):145-155
3. Egle Zasciurinskiene, Rune Lindsten, Christer Slotte, Krister Bjerklin. Orthodontic treatment in periodontitis-susceptible subjects: a systematic literature review. *Clinical and Experimental Dental Research*. 2016:162-173
4. Harun Canoglu, H. Cem Gungor, Zafer C. Cehreli. Management of cervical root fracture using orthodontic extrusion and crown reattachment: a case report. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 2007;104:e46-49
5. Elisabeth Reichardt, Ralf Krug, Michael M. Bornstein, Jürgen Tomasch, Carlalberta Verna, Gabriel Krastl. Orthodontic Forced Eruption of Permanent Anterior Teeth with Subgingival Fractures: A Systematic Review. *Int J Environ Res Public Health*. 2021,

18, 12580.

6. Grace Huang, Min Yang, Mohammad Qali, Tun-Jan Wang, Chenshuang Li, Yu-Cheng Chang. *Clinical Considerations in Orthodontically Forced Eruption for Restorative Purposes.* *J Clin Med.* 2021, 10, 5950.
7. Gaetano Isola, Riccardo Nucera, Silvia Damonte, Alessandro Ugolini, Anna De Mari, Marco Migliorati. *Implant Site Changes in Three Different Clinical Approaches: Orthodontic Extrusion, Regenerative Surgery and Spontaneous Healing after Extraction: A Systematic Review.* *J Clin Med.* 2022, 11, 6347.
8. BS Suprabha, M Kundabala, M Subraya, P Kancherla. *Reattachment and Orthodontic Extrusion in the management of an incisor crown-root fracture: A case report.* *J Clin Pediatr Dent* 30(3): 211-214, 2006.